arXiv:hep-ph/0212409v3 12 May 2003

The investigation of spin effects in photon production with fermion pair in $\gamma\gamma$-collisions

Author T.V. Shishkina

NC PHEP BSU
153 Bogdanovitcha str.,220040 Minsk, Belarus
E-mail: shishkina@hep.by

Author V.V. Makarenko

NC PHEP BSU
153 Bogdanovitcha str.,220040 Minsk, Belarus
E-mail: makarenko@hep.by

Abstract. The exclusive reaction $\gamma\gamma \to f\bar{f}\gamma$ is considered as the possible calibration process on the linear photon collider. We analyse the energy spectrum and total cross section dependence on the detector parameters at various initial beam helicities.

1. Introduction

Linear lepton colliders of the nearest future provide the possibility of investigation of photons collisions at energies and luminosities the same to those in e^+e^- collisions [1]. The photon linear colliders have the great physical potential (Higgs ans SUSY particles searching, study of anomalous gauge boson couplings and hadronic structure of photons etc., [2]). And performing of this set of investigations requires the of photon beam calibration as well. For this purpose one traditionally use some of the well-known and precisely calculated reactions (see, for example, $\gamma\gamma \to 2f, 4f$, [5, 6]). The difficulties appear in the calibration of photon beams of similar helicity due to the small magnitude of cross sections of the most QED processes.

For this purpose we consider the exclusive reaction $\gamma\gamma \to f\bar{f}\gamma$ as the possible candidate for a calibration channel on a linear photon collider.

The two various helicity configuration of the $\gamma\gamma$-system leads to the different spectra of final particles and requires the two mechanisms of beam calibration. We have analysed the behaviour of the $\gamma\gamma \to f\bar{f}\gamma$ reaction on beams with different helicities as a function of the parameters of detectors. We have performed the detail comparison of it's cross sections on $\gamma^+\gamma^+$- (total helicity $J = 0$) and $\gamma^+\gamma^-$-beams ($J = 2$). The question to be answered is does the process on ”++” beams can have the same cross section as one on ”+−” beams. We have outlined the conditions that greatly restrict the observation of the process on ”$J = 2$” beams, remaining the ”$J = 0$” cross section almost unchanged.



Fig. 1. Diagrams for the process $\gamma\gamma \to f\bar{f} + \gamma$.

We consider the process

$$\gamma(p_1, \lambda_1) + \gamma(p_2, \lambda_2) \to f({p_1}', {e_1}') + \bar{f}({p_2}', {e_2}') + \gamma(p_3, \lambda_3), \tag{1}$$

where λ_i and ${e_i}'$ are photon and fermion helicities.

We use the following set of invariants:

$$\begin{array}{lll}
s = (p_1 + p_2)^2 = 2\, p_1 \cdot p_2\,, & t = ({p_2}' - p_2)^2, & u = ({p_2}' - p_1)^2, \\
s' = ({p_1}' + {p_2}')^2, & t' = ({p_1}' - p_1)^2, & u' = ({p_1}' - p_2)^2, \\
\upsilon = 2\, {p_1}' \cdot p_3\,, & \nu = 2\, {p_2}' \cdot p_3\,, & \\
z = 2\, p_1 \cdot p_3\,, & z' = 2\, p_2 \cdot p_3\,. &
\end{array}$$

We denote the final-state photon energy by w, the polar angle (between any final and initial particles in c.m.s.) by Θ and the angle between any of final particles by φ. For the differential cross-section we introduce the normalized final-state photon energy (c.m.s. is used) $x = w/\sqrt{s}$. The differential cross section $d\sigma/dx$ appears to be the energy spectrum of final-state photons. The total helicity of the $\gamma\gamma$ system is denoted by J.

2. Calculation

We consider the cross section

$$\sigma = \int \frac{1}{2s} \left| M(\lambda_1, \lambda_2, {e_1}', {e_2}', \lambda_3) \right|^2 d\Gamma,$$

where the phase-space volume element is defined by

$$d\Gamma = \frac{d^3 p_1'}{(2\pi)^3\, 2\mathcal{E}_1'} \cdot \frac{d^3 p_2'}{(2\pi)^3\, 2\mathcal{E}_2'} \cdot \frac{d^3 p_3}{(2\pi)^3\, 2p_3^0} \cdot (2\pi)^4 \delta\left(p_1 + p_2 - p_1' - p_2' - p_3\right).$$

The process $\gamma\gamma \to f\bar{f}\gamma$ is the pure QED reaction in the Born approximation. Using the method of helicity amplitudes [3], we calculate matrix elements:

$$\left| M^{+--++} \right| = 2e^3\, p_2' \cdot p_2 \sqrt{\frac{p_1' \cdot p_2'}{p_1' \cdot p_3\ p_2' \cdot p_3\ p_1' \cdot p_1\ p_2' \cdot p_1}} = e^3 \sqrt{\frac{8st^2}{\nu \upsilon t' u}}. \tag{2}$$

All the other non-vanishing amplitudes are obtained from $|M^{+--++}|$ by using C, P, Bose and crossing (between final and initial particles) symmetries:

$$\begin{gathered}
d\sigma^{+-+--} = d\sigma^{+--++}|_{1\leftrightarrow 2},\ (P + Bose) \\
d\sigma^{+-+-+} = d\sigma^{+--++}|_{1'\leftrightarrow 2'},\ (C) \\
d\sigma^{+--+-} = d\sigma^{+--++}|_{\substack{1\leftrightarrow 2\\ 1'\leftrightarrow 2'}},\ (CP + Bose) \\
d\sigma^{+++--} = d\sigma^{+--++}|_{\substack{3\leftrightarrow 2\\ 1'\leftrightarrow 2'}},\ (C + crossing) \\
d\sigma^{++-+-} = d\sigma^{+++--}|_{1'\leftrightarrow 2'},\ (C) \\
d\sigma^{-\lambda_1, -\lambda_2, -{e_1}', -{e_2}', -\lambda_3} = d\sigma^{\lambda_1, \lambda_2, {e_1}', {e_2}', \lambda_3}.\ (P)
\end{gathered}$$

Due to no final-state polarizations can be measured we summarize over all the final particles helicities.

The further integration is performed numerically using the Monte-Carlo method [4].

3. Results

One of the main purposes of the linear photon collider is the s-channel of the Higgs boson production at energies about $\sqrt{s} = 120 GeV$ [7]. Hence we use this value of c.m.s. energy in our analysis.

We have performed calculations for various experimental restrictions on the parameters of final particles. The events are not detected if Θ, φ or w are below the threshold values. The default minimal values are $\Theta_{min} = 7^o$, $\varphi_{min} = 3^o$, $w_{min} = 1 GeV$, the minimal fermion energy was also assumed to be $1 GeV$.

First of all we discuss the energy spectrum of final photons. As it is presented in fig. 2-3, the differential cross section ($d\sigma/dx$) on J=2 beams decreases while one on J=0 beams raises with increasing of the final-state photon energy. This leads to the conclusion that if one raise the threshold on w, the process on J=2 beams will be greatly restricted, but the rate of J=0 events remains almost unchanged. In fig. 4 we present the total cross section dependence on the w-cut. According to fig. 4c, one can achive the ratio $\sigma_{J=0}/\sigma_{J=2}$ up to 0.5 without sufficient decreasing of $\sigma_{J=2}$.

Next we analyse the total cross section dependence on the angle cuts (Θ_{min} and φ_{min}, see figs. 5-6). In $\gamma^+\gamma^+$- experiments ($J = 0$) most of final fermions are radiated closely to the axis of initial photon (polar axis) and can't be detected. The $\gamma^+\gamma^-$- experiments ($J = 2$) have the large fraction of particles emitted at large angles. But they are likely emitted as the collinear fermion-photon pairs, that also can't be separated in the detector. In the figs. 5c and 6c we present the ratio of the total cross sections $\sigma_{J=0}/\sigma_{J=2}$ at certain configuration of cuts. We conclude that it can be increased by reducing the polar angle cut Θ_{min}, and by raising the collinear angle cut φ_{min}.

In fig. 7 we demonstrate the total cross section dependence on the c.m.s. energy. It is close to the law $1/s$ due to all the invariants in calculations are proportional to s. The only non-linear effect is the w-cut.

We discover that the ratio of events on J=0 and J=2 beams strongly depends on the experimental cuts. We obtained the region (the configuration of cuts) where the processes on the both J=0 and J=2 beams have the close cross sections. That is the region of small polar angle cut, high collinear angle cut and high minimal energy of final-state photons. At these parameters the total cross sections of $\gamma\gamma \to f\bar{f}\gamma$ in experiments using $\gamma^+\gamma^+$- and $\gamma^+\gamma^-$- beams appear to be the same order of magnitude.

References

[1] I.F. Ginzburg, G.L. Kotkin, V.G. Serbo and V.I. Telnov, Nucl. Instr. Meth. **205** (1983) 47;
I.F. Ginzburg, G.L. Kotkin, S.L. Panfil, V.G. Serbo and V.I. Telnov, Nucl. Instr. Meth. **219** (1984) 5.

[2] TESLA TDR Part 6. Chapter 1. Photon collider at TESLA, hep-ex/0108012;
E.A. Kuraev, M.V. Galynskii, M.I. Levchuk, Phys. Part. Nucl. **31** (2000) 76, Fiz. Elem. Chast. Atom. Yadra **31** (2000) 155;
V. Telnov, Nucl. Instr. Meth. **A494** (2002) 35, hep-ex/0207093.

[3] P. De Causmaecker, R. Gastmans, W. Troost, Tai Tsun Wu, Phys. Lett. **B105** (1981) 215;
P. De Causmaecker, R. Gastmans, W. Troost, Tai Tsun Wu, Nucl. Phys. **B206** (1982) 53;
F.A. Berends et al., Nucl. Phys. **B206** (1982) 61;
S. Dittmaier, Phys. Rev. **D59** (1999) 016007, hep-ph/9805445.

[4] S. Weinzierl, NIKHEF-00-012, hep-ph/0006269.

[5] A. Denner, S. Dittmaier, Eur. Phys. J. **C9** (1999) 425, hep-ph/9812411;
J. Layssac, F.M. Renard, Phys. Rev. **D64** (2001) 053018, hep-ph/0104205.

[6] C. Carimalo, W. da Silva, F. Kapusta, Nucl. Phys. Proc. Suppl. **82** (2000) 391, hep-ph/9909339;
M. Moretti, Nucl. Phys. **B484** (1997) 3, hep-ph/9604303.

[7] D.L. Borden, D.A. Bauer, D.O. Caldwell, Phys. Rev. **D48** (1993) 4018;
J.F. Gunion, H.E. Haber, Phys. Rev. **D48** (1993) 5109.



Fig. 2. The cross section $d\sigma/dx$ for $\sqrt{s} = 120GeV$. The cuts are: $5^o, 7^o$ and 10^o on the polar angle, 3^o on the collinear angle and $1GeV$ on the minimal final-state particle energy.



Fig. 3. The cross section $d\sigma/dx$ for $\sqrt{s} = 500GeV$. The cuts are: $5^o, 7^o$ and 10^o on the polar angle, 3^o on the collinear angle and $1GeV$ on the minimal final-state particle energy.



Fig. 4. Total cross section dependence on the final photon minimal energy. The cuts are: 7^o on the polar angle, 3^o between final particles and $1GeV$ on the minimal fermion energy.



Fig. 5. Total cross section dependence on the polar angle cut. The other cuts are: 3^o between final particles and $1GeV$ on the minimal final-state particle energy.



Fig. 6. Total cross section dependence on the minimal angle between final particles. The other cuts are: 7^o on the polar angle and $1 GeV$ on the minimal final-state particle energy.



Fig. 7. The dependence of total cross section on the c.m.s. energy. The cuts are: 7^o (polar angle), 3^o (collinear angle), $1 GeV$ (minimal fermion energy), $1 GeV$ (w_{cut}, the higher line) and $5 GeV$ (w_{cut}, the lower line).